

$$\mathcal{F}_{2\pi\text{-per}}(\mathbb{R},\mathbb{C}) = \left\{ f: \mathbb{R} \longrightarrow \mathbb{C} \mid f(x+2\pi) = f(x) \text{ for all } x \in \mathbb{R} \right\}$$

Let's take integrable functions:

$$\mathcal{L}_{2\pi\text{-per}}^{1}(\mathbb{R},\mathbb{C}) = \left\{ f \in \mathcal{F}_{2\pi\text{-per}}(\mathbb{R},\mathbb{C}) \mid \int_{1}^{\pi} |f(x)| \, dx < \infty \right\}$$

$$f \text{ integrable with respect to} Lebesgue measure on [-\pi, \pi]$$

$$\frac{\text{norm?}}{\|f\|_{1}} := \int_{\pi}^{\pi} |f(x)| \, dx \qquad \frac{\text{problem:}}{\int_{\pi}^{\pi}} \int_{\pi}^{\pi} |f(x)| \, dx \qquad \frac{\text{problem:}}{\int_{\pi}^{\pi}} \int_{\pi}^{\pi} \int_{\pi}^{\pi}$$

<u>solution</u>: equivalence relation $f \sim g : \iff \|f - g\|_{1} = 0$ set of all equivalence classes: $L_{2\pi-per}^{1}(\mathbb{R},\mathbb{C}) := \mathcal{I}_{2\pi-per}^{1}(\mathbb{R},\mathbb{C})/\mathcal{A}$ [

 $\left\| \begin{bmatrix} f \end{bmatrix} \right\|_{1} := \left\| f \right\|_{1}$ > norm!

complex vector space

 $\frac{\text{identify:}}{2^{2} r - per} (\mathbb{R}, \mathbb{C}) \supseteq \mathcal{P}_{2^{2} r - per} (\mathbb{R}, \mathbb{C})$

Let's take square-integrable functions:

$$\mathcal{L}_{2\pi-per}^{2}(\mathbb{R},\mathbb{C}) = \left\{ f \in \mathcal{F}_{2\pi-per}(\mathbb{R},\mathbb{C}) \mid \int_{-\pi}^{\pi} |f(x)|^{2} dx < \infty \right\}$$

$$\frac{\operatorname{norm}^{9}}{||f||_{2}} := \sqrt{\int_{-\pi}^{\pi} |f(x)|^{2} dx}$$

<u>solution</u>: equivalence relation $f \sim g : \iff \|f - g\|_{2} = 0$ set of all equivalence classes: $\int_{2\pi - per}^{2} (\mathbb{R}, \mathbb{C}) := L_{2\pi - per}^{2} (\mathbb{R}, \mathbb{C}) / \mathcal{A}$ \bigotimes complex vector space with inner product