

Functional analysis - part 25

Hahn-Banach theorem $(X, \|\cdot\|_X)$ normed space $\rightsquigarrow (X', \|\cdot\|_{X'})$

$U \subseteq X$ subspace, $u': U \rightarrow \mathbb{F}$ continuous linear functional

Then: There exists $x': X \rightarrow \mathbb{F}$ continuous linear functional

with $x'(u) = u'(u)$ for all $u \in U$,

$$\|x'\|_{X'} = \|u'\|_{U'}$$

Applications: $(X, \|\cdot\|_X)$ normed space

(a) For all $x \in X, x \neq 0$, there is an $x' \in X'$ with $\|x'\|_{X'} = 1$ and $x'(x) = \|x\|_X$.


Proof: Define $u': U \rightarrow \mathbb{F}$

$$\lambda \cdot x \mapsto \lambda \|x\|_X \quad \text{continuous linear functional}$$

Hahn-Banach

$$\Rightarrow x': X \rightarrow \mathbb{F} \quad \text{with} \quad x'(x) = u'(x) = \|x\|_X$$

$$\|x'\|_{X'} = \|u'\|_{U'} = 1$$



(b) X' separates the points of X : For $x_1, x_2 \in X, x_1 \neq x_2$, there is an $x' \in X'$ with $x'(x_1) \neq x'(x_2)$

Proof: $x := x_2 - x_1 \stackrel{(a)}{\Rightarrow} x'(x) = \|x\|_X \neq 0 \Rightarrow x'(x_1) \neq x'(x_2)$

(c) For all $x \in X$: $\|x\|_X = \sup \{ |x'(x)| \mid x' \in X', \|x'\| = 1 \}$

Proof:

$$\|x'\|_{X'} \geq \frac{|x'(x)|}{\|x\|_X} \Rightarrow 1 = \sup_{\|x'\|=1} \|x'\|_{X'} \geq \sup_{\|x'\|=1} \frac{|x'(x)|}{\|x\|_X}$$

$$\Rightarrow \|x\|_X \geq \sup_{\|x'\|=1} |x'(x)|$$

Use (a):

$$\|x\|_X \leq \sup_{\|x'\|=1} |x'(x)|$$

(d) Let $U \subseteq X$ be a closed subspace, $x \in X$ with $x \notin U$.

Then there exists $x' \in X'$ with $x'|_U = 0$ and $x'(x) \neq 0$.

Proof: $X/U := \{ [z] \mid z \in X \}, [z] := \{ z + u \mid u \in U \}$

$$\|[z]\|_{X/U} := \inf_{u \in U} \|z + u\|_X \rightsquigarrow (X/U, \|\cdot\|_{X/U}) \text{ normed space}$$

$\stackrel{(a)}{\Rightarrow}$ There is a $\gamma' \in (X/U)'$ with $\gamma'([x]) \neq 0$.

Define $x' \in X'$ by $x'(z) := \gamma'([z])$ for $z \in X$.

