

## Functional analysis – part 31

Spectral radius:  $X$  complex Banach space  $T: X \rightarrow X$  bounded linear operator

$r(T) := \sup_{\lambda \in \sigma(T)} |\lambda|$

Theorem:  $X$  complex Banach space,  $T: X \rightarrow X$  bounded linear operator.

Then: (a)  $\sigma(T) \subseteq \mathbb{C}$  is compact

(b)  $X \neq \{0\} \Rightarrow \sigma(T) \neq \emptyset$

(c)  $\Gamma(T) := \sup_{\lambda \in \sigma(T)} |\lambda| = \lim_{k \rightarrow \infty} \|T^k\|^{\frac{1}{k}} = \inf_{k \in \mathbb{N}} \|T^k\|^{\frac{1}{k}} \leq \|T\| < \infty$

Proof: For  $\lambda \in \mathbb{C}$  with  $|\lambda| > \|T\|$ :  $(I - \frac{T}{\lambda})^{-1} = \sum_{k=0}^{\infty} \left(\frac{T}{\lambda}\right)^k$

$$\Rightarrow (T - \lambda)^{-1} = -\frac{1}{\lambda} (I - \frac{T}{\lambda})^{-1} = -\frac{1}{\lambda} \sum_{k=0}^{\infty} \left(\frac{T}{\lambda}\right)^k \quad (*)$$

$\Rightarrow \sup_{\lambda \in \sigma(T)} |\lambda| \leq \|T\| \Rightarrow \sigma(T)$  is bounded

For (b): Assume  $\sigma(T) = \emptyset \Rightarrow \rho(T) = \mathbb{C}$

Reminder: The map  $\rho(T) \rightarrow \mathcal{B}(X)$

$$\lambda \mapsto (T - \lambda)^{-1} \text{ is analytic.}$$

$$\text{Take any } \ell \in \mathcal{B}(X)^*: f_\ell: \mathbb{C} \rightarrow \mathbb{C}$$

$$\lambda \mapsto \ell((T - \lambda)^{-1})$$

analytic function (holomorphic function)

We get that  $f_\ell$  is a bounded entire function.

$$\hookrightarrow \text{For } |\lambda| \geq 2\cdot \|T\| : (T - \lambda)^{-1} = -\frac{1}{\lambda} \sum_{k=0}^{\infty} \left(\frac{T}{\lambda}\right)^k \quad (*)$$

$$|f_\ell(\lambda)| \leq \|\ell\| \cdot \|(T - \lambda)^{-1}\| \leq \|\ell\| \underbrace{\frac{1}{|\lambda|}}_{\leq \frac{1}{2\|T\|}} \underbrace{\sum_{k=0}^{\infty} \left\| \frac{T}{\lambda} \right\|^k}_{\leq \frac{1}{2}}$$

Liouville's theorem

$$\implies f_\ell \text{ is constant}$$

$$f_\ell(0) = \ell(T^{-1})$$

$$\begin{aligned} f_\ell(\lambda) &= \ell((T - \lambda)^{-1}) = \ell\left(\sum_{k=0}^{\infty} (T)^{-(k+1)} \cdot (\lambda)^k\right) \\ &= \sum_{k=0}^{\infty} \ell(T^{-(k+1)}) \cdot \lambda^k \end{aligned}$$

$$\implies \ell(T^{-1}) = 0 \text{ for all } \ell \in \mathcal{B}(X)^*$$

Hahn-Banach theorem

$$\implies_{(\text{part 25})} T^{-1} = 0 \implies X = \{0\}$$