


Ordinary Differential Equations – Part 3

ODE: $\dot{x} = w(t, x)$ (explicit, of first order)


- Example:
- (a) $\dot{x} = \lambda \cdot x \rightsquigarrow$ autonomous
 - (b) $\dot{x} = t \rightsquigarrow$ not autonomous
 - (c) $\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ -x_1 \end{pmatrix} \rightsquigarrow$ autonomous

Definition: autonomous system: $\dot{x} = v(x)$ with $v: U \rightarrow \mathbb{R}^n$ often:
 U open
 v continuous

Directional field:



Examples: (a) $\dot{x} = \sin(x)$, $v: \mathbb{R} \rightarrow \mathbb{R}$, $v(x) = \sin(x)$



(1) $\alpha(t) = 0$ for all $t \in \mathbb{R}$ is a solution: $\dot{\alpha}(t) = \underbrace{\sin(\alpha(t))}_{=0} = 0$

(2) $\alpha(t) = \pi$ for all $t \in \mathbb{R}$ is a solution.

(3) A solution with $\alpha(0) = \frac{\pi}{2}$ is monotonically increasing
with $\lim_{t \rightarrow \infty} \alpha(t) = \pi$.

$$(b) \quad \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}, \quad \forall: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x_1, x_2) \mapsto \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}$$

