

Real Analysis - Part 27


Definition: Let $f: I \rightarrow \mathbb{R}$ be a function with $I \subseteq \mathbb{R}$.

f is called continuous at $x_0 \in I$ if

$$\lim_{x \rightarrow x_0} f(x) = f(x_0)$$

or if x_0 is isolated in I .

There is no sequence $(x_n)_{n \in \mathbb{N}} \subseteq I \setminus \{x_0\}$ with $\lim_{n \rightarrow \infty} x_n = x_0$




Definition: Let $f: I \rightarrow \mathbb{R}$ be a function with $I \subseteq \mathbb{R}$.


f is called continuous (on I) if f is continuous at x_0 for all $x_0 \in I$.

To remember: Continuity implies: $\lim_{n \rightarrow \infty} f(x_n) = f\left(\lim_{n \rightarrow \infty} x_n\right)$ (if $\lim_{n \rightarrow \infty} x_n \in I$)


Examples: (a) $f: I \rightarrow \mathbb{R}$ constant



$$(b) f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = \begin{cases} 0 & , x < 0 \\ 1 & , x \geq 0 \end{cases}$$



$$(c) f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = \begin{cases} 0 & , x \neq 0 \\ 1 & , x = 0 \end{cases}$$




(d) $f: \mathbb{R} \rightarrow \mathbb{R}$ polynomial

$$f(x) = a_m \cdot x^m + a_{m-1} \cdot x^{m-1} + \dots + a_1 \cdot x^1 + a_0$$

We have: $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ for all $x_0 \in I$.

limit theorem for sequences



(e) $f: I \rightarrow \mathbb{R}$ rational function


$$I := \{x \in \mathbb{R} \mid q(x) \neq 0\}$$

$$f(x) = \frac{\overset{\text{polynomial}}{p(x)}}{\underset{\text{polynomial}}{q(x)}}$$

continuous on I

(f) $f: \mathbb{R} \rightarrow \mathbb{R}$ absolute value

$$f(x) = |x| = \begin{cases} -x & , x < 0 \\ x & , x \geq 0 \end{cases}$$



$$\left. \begin{aligned} \lim_{x \rightarrow 0} f(x) &= \lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} (-x_n) = 0 \\ \lim_{x \rightarrow 0} f(x) &= \lim_{n \rightarrow \infty} f(x_n) = \lim_{n \rightarrow \infty} (x_n) = 0 \end{aligned} \right\} \Rightarrow \lim_{x \rightarrow x_0} f(x) = f(x_0)$$

(g) $f: \mathbb{R} \rightarrow \mathbb{R}$ $f(x) = \begin{cases} 0 & , x \notin \mathbb{Q} \\ 1 & , x \in \mathbb{Q} \end{cases}$

(\mathbb{Q} is dense in \mathbb{R} by construction)

