


## Real Analysis - Part 40

Definition:  $I \subseteq \mathbb{R}$  interval,  $f: I \rightarrow \mathbb{R}$ .



(a)  $f$  has a local maximum at  $x_0 \in I$  if there is a neighbourhood of  $x_0$ ,  $U \subseteq \mathbb{R}$ , with

$$f(x_0) = \max \{ f(x) \mid x \in U \cap I \}$$


(b)  $f$  has a local minimum at  $x_0 \in I$  if there is a neighbourhood of  $x_0$ ,  $U \subseteq \mathbb{R}$ , with



$$f(x_0) = \min \{ f(x) \mid x \in U \cap I \}$$

(c)  $f$  has a local extremum at  $x_0 \in I$  if  $f$  has a local maximum or local minimum at  $x_0 \in I$ .



Proposition:  $f: (a, b) \rightarrow \mathbb{R}$  differentiable at  $x_0 \in (a, b)$ .

$f$  has a local extremum at  $x_0 \Rightarrow f'(x_0) = 0$




Proof: 1st case:  $f$  has a local maximum at  $x_0$

$\Rightarrow$  there is a neighbourhood of  $x_0$ ,  $U \subseteq (a, b)$


$$f(x_0) = \max \{ f(x) \mid x \in U \}$$

$$f \text{ differentiable at } x_0 \Rightarrow f(x) = f(x_0) + (x - x_0) \cdot \underbrace{\Delta_{f, x_0}(x)}_{\text{continuous at } x_0}$$



Assume  $f'(x_0) > 0$ : There exists a neighbourhood  $V \subseteq U$  such that  $\Delta_{f, x_0}(x) > 0$  for all  $x \in V$ .

$$\text{Then: } x > x_0 \Rightarrow f(x) = f(x_0) + \underbrace{(x - x_0)}_{> 0} \cdot \underbrace{\Delta_{f, x_0}(x)}_{> 0} > f(x_0)$$



Assume  $f'(x_0) < 0$ : There exists a neighbourhood  $V \subseteq U$  such that  $\Delta_{f, x_0}(x) < 0$  for all  $x \in V$ .

$$\text{Then: } x < x_0 \Rightarrow f(x) = f(x_0) + \underbrace{(x - x_0)}_{< 0} \cdot \underbrace{\Delta_{f, x_0}(x)}_{< 0} > f(x_0)$$


$$\Rightarrow f'(x_0) = 0$$

2nd case:  $f$  has a local minimum at  $x_0$  (works similarly)

### Theorem of Rolle

$f: [a, b] \rightarrow \mathbb{R}$  differentiable and  $f(a) = f(b)$ .

Then there is  $\hat{x} \in (a, b)$  with  $f'(\hat{x}) = 0$ .



Proof: 1st case:  $f$  constant  $\Rightarrow f'(x) = 0$  for all  $x \in [a, b]$ . ✓

2nd case:  $f$  is not constant.

There are  $x^-, x^+ \in [a, b]$  with  $f(x^+) = \sup\{f(x) \mid x \in [a, b]\}$   
 $f(x^-) = \inf\{f(x) \mid x \in [a, b]\}$

$f$  not constant

$\Rightarrow x^- \in (a, b)$  or  $x^+ \in (a, b)$  (call it  $\hat{x}$ )

Proposition above

$\Rightarrow f'(\hat{x}) = 0$

□