


## Start Learning Reals - Part 1

Real numbers  $\mathbb{R}$

Starting point:  $\mathbb{Q}$  is the set of fractions  $\rightsquigarrow$  field and Archimedean order  $\leq$   
 $x > 0$  ,  $x < 0$

Absolute value: For  $x \in \mathbb{Q}$  define:  $|x| := \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$



How far away is  $x$  from  $0$ ?  $\rightsquigarrow |x|$

Problem: There is no  $x \in \mathbb{Q}$  with  $x^2 = 2$

$$x_1 = \frac{14}{10} = \frac{7}{5} \rightsquigarrow x_1^2 = \frac{49}{25} \approx 2$$

$$x_2 = \frac{141}{100} \rightsquigarrow x_2^2 = \frac{19881}{10000} \approx 2$$

$$x_3 = \frac{1414}{1000} \rightsquigarrow x_3^2 = \frac{499849}{250000} \approx 2$$

$$x_4 = \frac{14142}{10000} \rightsquigarrow x_4^2 = \frac{4999041}{25000000} \approx 2$$

$$x_5 = \frac{141421}{100000} \rightsquigarrow x_5^2 = \frac{1999899241}{10000000000} \approx 2$$

$\vdots$   $\vdots$   $\vdots$

$$x = ? \rightsquigarrow x^2 = 2$$


distance:

$$|x_5 - x_3|$$

We consider a sequence  $(x_n)_{n \in \mathbb{N}}$  (infinite list; formally: a map  $\mathbb{N} \rightarrow \mathbb{Q}$ ,  $n \mapsto x_n$ )  
 with the property:

$$\forall \varepsilon \in \mathbb{Q} \exists N \in \mathbb{N} \forall n, m \in \mathbb{N} : (\varepsilon > 0 \wedge n, m \geq N \implies |x_n - x_m| < \varepsilon)$$

In short:  $\forall \varepsilon > 0 \exists N \in \mathbb{N} \forall n, m \geq N : |x_n - x_m| < \varepsilon \quad (*)$



Cauchy sequence: sequence  $(x_n)_{n \in \mathbb{N}}$  with  $x_n \in \mathbb{Q}$  and property  $(*)$